

ON THE INTERRELATIONSHIPS AMONG SUFFICIENCY,  
TOTAL SUFFICIENCY AND SOME RELATED CONCEPTS

BY

STEFFEN L. LAURITZEN

TECHNICAL REPORT NO. 18

JUNE, 1974

PREPARED UNDER CONTRACT

N00014-67-A-0112-0030 (NR-042-034)

FOR THE OFFICE OF NAVAL RESEARCH

THEODORE W. ANDERSON, PROJECT DIRECTOR

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA



ON THE INTERRELATIONSHIPS AMONG SUFFICIENCY,  
TOTAL SUFFICIENCY AND SOME RELATED CONCEPTS

by

STEFFEN L. LAURITZEN  
University of Copenhagen

and

Stanford University

TECHNICAL REPORT NO. 18

JUNE, 1974

PREPARED UNDER THE AUSPICES

OF

OFFICE OF NAVAL RESEARCH CONTRACT #N00014-67-A-0112-0030

(NR-042-034)

THEODORE W. ANDERSON, PROJECT DIRECTOR

DEPARTMENT OF STATISTICS

STANFORD UNIVERSITY

STANFORD, CALIFORNIA

## 1. Introduction.

The literature on sufficiency is extensive and it is not the aim of the present paper to give a complete survey of this. We shall discuss the relationship between a number of notions introduced by various authors with different problems in mind, but all of them being of the same nature as sufficiency. Some of these notions were defined in terms of subfields of abstract probability spaces, but we shall restate all definitions in terms of statistics and discrete probabilities as our interest is directed more towards structural properties than technical ones.

## 2. Sufficiency, adequacy and summarizing statistics.

In the present section we shall investigate three different properties of statistics with the same basic idea, namely that they express the intuitive statement that a statistic contains all "relevant" information.

First the classical notion of a sufficient statistic as introduced by Fisher (1920). We shall define it the following way:

Let  $X$  be a random variable on a discrete, at most denumerable space  $E$  and  $t$  a mapping from  $E$  into another discrete space  $F$ . Let  $\mathcal{P}$  be a family of probabilities on  $E$  and let  $Y = t(X)$ .

Definition 2.1.  $t$  is said to be sufficient for  $\mathcal{P}$  if there is a fixed non negative real function  $\varphi$  on  $E \times F$  so that for all  $P \in \mathcal{P}$  and all  $x \in E$ :

$$\begin{aligned} P\{Y = y\} &> 0 \\ \Rightarrow P\{X = x | Y = y\} &= \varphi(x, y). \end{aligned}$$

A slightly stronger notion was introduced by Freedman (1962) with the pure probabilistic motivation of generalizing de Finetti's theorem for exchangeable 0-1 random variables. The notion is however closely related to sufficiency, as we shall soon see. Again let  $X$  be a random variable on a discrete space  $E$  and  $t$  a mapping from  $E$  to a discrete space  $F$ .

Definition 2.2. A probability measure  $P$  on  $E$  is said to be summarized by  $t$  if for all  $x, x' \in E$

$$t(x) = t(x') \Rightarrow P\{X = x\} = P\{X = x'\}.$$

In contrast to definition 2.1, we are not dealing with a family of probabilities but only with one probability measure. To be able to see the relation between a sufficient statistic and a summarizing statistic we have to define a summarizing statistic for a family of probabilities  $\mathcal{P}$ . In the previous notation we define

Definition 2.3.  $t$  is said to summarize  $\mathcal{P}$  if all  $P \in \mathcal{P}$  are summarized by  $t$ .

Remark: Note that the term "summarizing" is essentially related to discrete random variables as opposed to other concepts dealt with in the present paper.

This is stronger than sufficiency:

Proposition 2.1. If  $\mathcal{P}$  is a family of probability measures on  $E$  and  $t: E \rightarrow F$  summarizes  $\mathcal{P}$ , then  $t$  is sufficient for  $\mathcal{P}$ .

Proof. We shall just specify the function  $\varphi$  in definition 1.

Define  $\varphi$  as

$$(1) \quad \varphi(x, y) = \begin{cases} 0 & \text{if } t(x) \neq y \\ \frac{1}{N(y)} & \text{for } t(x) = y, \end{cases}$$

where  $N(y)$  is the total number of  $x$ 's so that  $t(x) = y$ ,  $0 < N(y) \leq \infty$ .

If  $P\{Y = y\} > 0$ , we have

$$(2) \quad \begin{aligned} P\{X = x | Y = y\} \\ = \frac{P\{X = x \wedge Y = y\}}{P\{Y = y\}} = x_{t^{-1}(y)}(x) \cdot \frac{P\{X = x\}}{\sum_{z \in t^{-1}(y)} P\{X = z\}}, \end{aligned}$$

where  $x_A$  is the indicator function of the set  $A$ :

$$x_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A. \end{cases}$$

If  $t(x) = y$ , we have

$$(3) \quad P\{X = z\} = P\{X = x\} \quad \text{for all } z \in t^{-1}(y),$$

since  $t$  was a summarizing statistic. Hence

$$\begin{aligned}
 (4) \quad P\{Y = y\} &= \sum_{z \in t^{-1}(y)} P\{X = z\} \\
 &= P\{X = x\} \cdot N(y) .
 \end{aligned}$$

So

$$(5) \quad P\{Y = y\} > 0 \implies N(y) < \infty$$

and

$$\begin{aligned}
 (6) \quad P\{X = x | Y = y\} &= x_{t^{-1}(y)}(x) \cdot \frac{P\{X = x\}}{P\{X = x\}N(y)} \\
 &= x_{t^{-1}(y)}(x) \cdot \frac{1}{N(y)} = \varphi(x, y) ,
 \end{aligned}$$

which was to be proved.

So the notion of a summarizing statistic is stronger than that of a sufficient statistic in the sense that not only is the conditional distribution of  $X$  given  $t(X)$  supposed to be known, but this distribution is supposed to have the specific form (6), i.e. uniform on the set  $t^{-1}(t(X))$ .

Barndorff-Nielsen and Skibinsky (1963) considered the problem of how much one could reduce a data set and still have all relevant information for the prediction of an unobserved random variable when the joint distribution of the data and the unobserved random variable was completely known and defined the notion adequacy. This definition was later extended to the case, where this joint distribution was only known to be a member of a specified family of distributions by Skibinsky (1967).

Let  $X$  and  $Z$  be random variables on discrete, at most denumerable spaces  $E$  and  $G$ . Let  $\mathcal{P}$  be a family of distributions on  $E \times G$  and let  $\mathcal{P}_E$  denote the induced family of marginal distributions of  $X$ . Let  $t$  be a mapping from  $E$  to an at most denumerable space  $F$ .

Definition 2.4.  $t$  is said to be adequate for  $Z$  if

- i)  $t$  is sufficient for  $\mathcal{P}_E$
- ii) for all  $P \in \mathcal{P}$ :  $P\{X = x\} > 0$   
 $\Rightarrow P\{Z = z | X = x\} = P\{Z = z | t(X) = t(x)\}$ .

This definition suggests that the classical notion of sufficiency is not satisfactory to the theory of statistical inference in stochastic processes as the prediction of unobserved random variables (the future of the process observed) in most cases will be relevant. In the next section we shall consider some extra conditions that have been imposed on a sequence of statistics by various authors.

### 3. Sequences of statistics.

In the present section we shall let  $X_1, X_2, \dots$  be a sequence of random variables on discrete at most denumerable spaces  $E_1, E_2, \dots$  and let  $\mathcal{P}$  be a family of probability measures on  $E_1 \times E_2 \times \dots$ . Let  $\mathcal{P}^{(n)}$  denote the family of marginal distributions of  $X_1, \dots, X_n$  induced by  $\mathcal{P}$ . We shall consider a sequence  $t_1, t_2, \dots$  of mappings.

$$t_n : E_1 \times \dots \times E_n \rightarrow F_n,$$

where  $F_n$  are discrete, at most denumerable, and let  $Y_n = t_n(X_1, \dots, X_n)$ .

Bahadur (1954) introduced the term of a sufficient and transitive sequence of statistics in connection with sequential decision theory, which can be stated as follows.

Definition 3.1. The sequence  $t_1, t_2, \dots$  is said to be sufficient and transitive if for all  $n$ ,  $t_n$  is adequate for  $Y_{n+1}$ .

In other words,  $t_1, t_2, \dots$  is sufficient and transitive iff it at each step  $n$  contains all information relevant for the prediction of the value of the next statistic. This is related to but different from the notion of a totally sufficient statistic, introduced by Lauritzen (1972) in terms of abstract measure spaces and restated in terms of discrete probability spaces in Lauritzen (1974).

Definition 3.2.  $t_n$  is said to be totally sufficient if it is adequate for  $X_{n+1}, \dots, X_{n+k}$  for all  $k = 1, 2, \dots$ .

That the two notions are different can be seen by the following example:

Example 1. Let  $X_1, X_2$  be independent Poisson distributed with mean  $\lambda > 0$ , and let  $X_n = X_2 + Z_1 + \dots + Z_{n-2}$  for  $n \geq 3$ , where  $Z_1, Z_2, \dots$  are independent of  $X_1, X_2$  and independent identically Poisson distributed with mean 1. The sequence  $t_1, t_2, \dots$  of mappings defined by

$$(7) \quad \begin{aligned} t_1(x) &= x \quad \text{and} \\ t_n(x_1, \dots, x_n) &= x_1 + x_2 \quad \text{for } n \geq 2 \end{aligned}$$

is sufficient and transitive, whereas e.g.  $t_2$  is not totally sufficient as  $(X_1, X_2)$  and  $X_3 = X_2 + X_1$  are not conditionally independent.



On the other hand, the sequence  $s_1, s_2, \dots$  defined as

$$(8) \quad \left\{ \begin{array}{l} s_1(x) = x \\ s_2(x_1, x_2) = (x_1, x_2) \\ s_3(x_1, x_2, x_3) = (x_1, x_2, x_3) \\ s_4(x_1, x_2, x_3, x_4) = (x_1, x_2, x_4) \\ s_5(x_1, x_2, x_3, x_4, x_5) = (x_1, x_2, x_3, x_5) \text{ and} \\ s_n(x_1, \dots, x_n) = (x_1, x_2, x_n) \text{ for } n \geq 6 \end{array} \right.$$

is totally sufficient but not sufficient and transitive, because

$(X_1, X_2, X_3, X_4)$  and  $s_5(X_1, \dots, X_5) = (X_1, X_2, X_3, X_5)$  are not conditionally independent given  $s_4(X_1, \dots, X_4) = (X_1, X_2, X_4)$ .

If one wants to insure a sequence of totally sufficient statistics to be sufficient and transitive, an extra condition has to be imposed.

The following algebraic property of a sequence of statistics is a slight weakening of "S-structure" as defined by Freedman (1962).

Definition 3.3.  $t_1, t_2, \dots$  is said to have  $\sum$ -structure if for all  $m, n$

$$t_n(x_1, \dots, x_n) = t_n(y_1, \dots, y_n)$$

$$\Rightarrow t_{n+m}(x_1, \dots, x_n, x_{n+1}, \dots, x_{n+m}) = t_{n+m}(y_1, \dots, y_n, x_{n+1}, \dots, x_{n+m})$$

In other words,  $t_1, t_2, \dots$  has  $\sum$ -structure if for all  $m$  and  $n$ , there is a mapping  $\psi_{nm}$

$$\psi_{nm}: F_n \times E_{n+1} \times \dots \times E_{n+m} \rightarrow F_{n+m}$$

so that

$$t_{n+m}(x_1, \dots, x_{n+m}) = \psi_{nm}(t_n(x_1, \dots, x_n), x_{n+1}, \dots, x_{n+m}) .$$

The term  $\sum$ -structure is chosen to emphasise that  $t_{n+m}$  is a "generalized sum" of  $t_n(x_1, \dots, x_n)$  and  $(x_{n+1}, \dots, x_{n+m})$ , as is e.g. the case in classical exponential families, where we have

$$t_n(x_1, \dots, x_n) = t(x_1) + \dots + t(x_n)$$

for  $t$  being some function from  $E$  into  $k$ -dimensional Euclidean space.

We can now show the following result:

Proposition 3.1. If for any  $n$ ,  $t_n$  is totally sufficient and if  $t_1, t_2, \dots$  has  $\sum$ -structure, then  $t_1, t_2, \dots$  is sufficient and transitive.

Proof.  $t_n$  is clearly sufficient for  $\mathcal{P}^{(n)}$  for all  $n$ . We have to show that  $Y_{n+1}$  and  $X_1, \dots, X_n$  are conditionally independent given  $Y_n$ . We get

$$\begin{aligned} & P\{Y_{n+1} = y | X_1 = x_1, \dots, X_n = x_n\} \\ (9) \quad & = P\{t_{n+1}(x_1, \dots, x_n, X_{n+1}) = y | X_1 = x_1, \dots, X_n = x_n\} \\ & = P\{\psi_{n,n+1}(t_n(x_1, \dots, x_n), X_{n+1}) = y | X_1 = x_1, \dots, X_n = x_n\} , \end{aligned}$$

where  $\psi_{n,n+1}$  satisfies

$$(10) \quad t_{n+1}(x_1, \dots, x_{n+1}) = \psi_{n,n+1}(t_n(x_1, \dots, x_n), x_{n+1}) .$$

As  $t_n$  is totally sufficient,  $X_{n+1}$  and  $X_1, \dots, X_n$  are conditionally independent given  $Y_n$ , and we get from (9) that

$$\begin{aligned}
 & P\{Y_{n+1} = y | X_1 = x_1, \dots, X_n = x_n\} \\
 (11) \quad & = P\{\psi_{n+1}(t_n(x_1, \dots, x_n), X_{n+1}) = y | Y_n = t_n(x_1, \dots, x_n)\} \\
 & = P\{Y_{n+1} = y | Y_n = t_n(x_1, \dots, x_n)\},
 \end{aligned}$$

which was to be proved.

Martin-Löf (1973) defined an algebraic consistency condition for a sequence of statistics which is slightly weaker than  $\Sigma$ -structure. We shall call it  $\Sigma^*$ -structure:

Definition 3.4.  $t_1, t_2, \dots$  is said to have  $\Sigma^*$ -structure if for all  $n, m$  there is a function

$$N_{n,m}: F_n \times F_m \rightarrow \{0, 1, \dots\}$$

so that for all  $(x_1, \dots, x_n) \in E_1 \times \dots \times E_n$

$$\begin{aligned}
 & \# \{ (x_{n+1}, \dots, x_{n+m}) \in E_{n+1} \times \dots \times E_{n+m} : t_{n+m}(x_1, \dots, x_{n+m}) = y \} \\
 & = N_{n,m}(t_n(x_1, \dots, x_n), y).
 \end{aligned}$$

Remark: The discreteness of the sample space is also essential here, as the number of points corresponding to given values of the statistics occur in the definition in a fundamental manner.

It is immediate that we have

Proposition 3.2. If  $t_1, t_2, \dots$  has  $\Sigma$ -structure, then it has  $\Sigma^*$ -structure.

Proof. Trivial.

$\sum^*$ -structure becomes an important property when the conditional distributions given the statistics are determined by the numbers

$$(12) \quad N_n(y) = \# \{ (x_1, \dots, x_n) \in E_1 \times \dots \times E_n : t_n(x_1, \dots, x_n) = y \}$$

and we have

Proposition 3.3. If for all  $n$ ,  $\mathcal{P}^{(n)}$  is summarized by  $t_n$  and if  $t_1, t_2, \dots$  has  $\sum^*$ -structure, then  $t_1, t_2, \dots$  is sufficient and transitive.

Proof. We already know that  $t_n$  is sufficient for  $\mathcal{P}^{(n)}$  from proposition 2.1. As in proposition 3.1 it remains to be shown that  $Y_{n+1}$  and  $X_1, \dots, X_n$  are conditionally independent given  $Y_n$ . We have for all  $n$

$$(13) \quad \begin{aligned} P\{X_1 = x_1, \dots, X_n = x_n\} \\ &= P\{X_1 = x_1, \dots, X_n = x_n | Y_n = t_n(x_1, \dots, x_n)\} \cdot P\{Y_n = t_n(x_1, \dots, x_n)\} \\ &= \frac{P\{Y_n = t_n(x_1, \dots, x_n)\}}{N_n(t_n(x_1, \dots, x_n))}, \end{aligned}$$

according to proposition 2.1.

Furthermore

$$(14) \quad \begin{aligned} P\{Y_{n+1} = y \wedge X_1 = x_1, \dots, X_n = x_n\} \\ &= \sum_{x: (x_1, \dots, x_n, x) \in t_{n+1}^{-1}(y)} \frac{P\{Y_{n+1} = y\}}{N_{n+1}(y)} \\ &= N_{n,n+1}(t_n(x_1, \dots, x_n)) \cdot \frac{P\{Y_{n+1} = y\}}{N_{n+1}(y)}. \end{aligned}$$

Hence from (13) and (14) we get

$$(15) \quad P\{Y_{n+1} = y | X_1 = x_1, \dots, X_n = x_n\} = \frac{P\{Y_{n+1} = y\} N_n(t_n(x_1, \dots, x_n))}{N_{n+1}(y) P\{Y_n = t_n(x_1, \dots, x_n)\}} N_{n,n+1}(t_n(x_1, \dots, x_n)).$$

Since this only depends on  $x_1, \dots, x_n$  through  $t_n(x_1, \dots, x_n)$ , we must have

$$(16) \quad \begin{aligned} P\{Y_{n+1} = y | X_1 = x_1, \dots, X_n = x_n\} \\ = P\{Y_{n+1} = y | Y_n = t_n(x_1, \dots, x_n)\}, \end{aligned}$$

which was to be proved.

If we assume  $\sum$ -structure instead of  $\sum^*$ -structure, we have the even stronger result:

Proposition 3.4. If for all  $n$ ,  $\mathcal{P}$  is summarized by  $t_n$  and if  $t_1, t_2, \dots$  has  $\sum$ -structure, then  $t_n$  is totally sufficient for all  $n$ .

Proof. As in the previous proof, we only have to establish that  $X_1, \dots, X_n$  and  $X_{n+1}, \dots, X_{n+k}$  are conditionally independent given  $Y_n$  for all  $n$  and  $k$ . We have

$$\begin{aligned}
& P\{X_1 = x_1, \dots, X_{n+k} = x_{n+k} | Y_n = y\} \\
&= \frac{x_{t_n^{-1}(y)}(x_1, \dots, x_n) \cdot P\{Y_{n+k} = t_{n+k}(x_1, \dots, x_{n+k})\}}{P\{Y_n = y\} \cdot N_{n+k}(t_{n+k}(x_1, \dots, x_{n+k}))} \\
(17) \quad &= \frac{x_{t_n^{-1}(y)}(x_1, \dots, x_n) \cdot P\{Y_{n+k} = \psi_{n,k}(y, x_{n+1}, \dots, x_{n+k})\}}{P\{Y_n = y\} \cdot N_{n+k}(\psi_{n,k}(y, x_{n+1}, \dots, x_{n+k}))},
\end{aligned}$$

where  $\psi_{n,k}$  is given by

$$(18) \quad \psi_{n,k}(t_n(x_1, \dots, x_n), x_{n+1}, \dots, x_{n+k}) = t_{n+k}(x_1, \dots, x_{n+k}).$$


But for fixed  $y$ , the expression (17) is a product in  $(x_1, \dots, x_n)$  and  $(x_{n+1}, \dots, x_{n+k})$ . Hence  $X_1, \dots, X_n$  and  $X_{n+1}, \dots, X_{n+k}$  are conditionally independent given  $Y_n = y$ , which was to be proved.

Apparently the property of being summarizing with  $\sum$ -structure is very strong. Another way of strengthening total sufficiency is to assume minimality. As in Lauritzen (1974) we say that  $t_n$  is minimal totally sufficient if it is a function of any other totally sufficient statistic. We then have

Proposition 3.5. If for all  $n$ ,  $t_n$  is minimal totally sufficient then  $t_1, t_2, \dots$  has  $\sum$ -structure.

Proof. The result follows directly from corollary 1 of Lauritzen (1974).

If we include the notion of a minimal sufficient statistic in our considerations, we can "summarize" the results in the following diagram:



(The implications that are not proved in the previous are trivial).

At this point the author feels uncomfortable as a statistician. Is it really so that all these various notions are relevant? It is certainly true that in many examples at least some of the notions coincide. So far we have dealt with  $\mathcal{P}$  being an arbitrary family of probability measures which in some sense is unreasonable from a statistical point of view. In the last section we shall impose regularity conditions on  $\mathcal{P}$  and see how many of the implications in the diagram turn into equivalences.

#### 4. Independence and universality.

Let us assume that for all  $P \in \mathcal{P}$ ,  $X_1, X_2, \dots$  are independent random variables. It is then immediate that total sufficiency and sufficiency coincide and the same is of course true for minimal total sufficiency and minimal sufficiency. Hence it appears from the diagram that e.g. "minimal sufficiency" implies everything but "summarizing" and is thus a very strong property.

Barndorff-Nielsen (1973) discussed the notion of a universal family of probability measures in connection with the notion of M-ancillarity. Let  $X$  be a random variable on a discrete, at most denumerable set  $E$  and  $\mathcal{P}$  a family of probability measures on  $E$ .

Definition 4.1.  $\mathcal{P}$  is said to be universal if for all  $x \in E$  there is a  $P \in \mathcal{P}$  so that

$$P\{X = x\} \geq P\{X = y\}$$

for all  $y \in E$ .

The following result, given in e.g. Barndorff-Nielsen (1973) shows a relation to the discussion in the preceding sections:

Proposition 4.1. If  $\mathcal{P}$  is universal and  $t$  is sufficient for  $\mathcal{P}$ , then  $t$  summarizes  $\mathcal{P}$ .

Proof. The proof is exactly as in Barndorff-Nielsen (1973), theorem 2.1. Although  $E$  is assumed to be finite in that paper, this assumption is irrelevant for the validity of the proof.

Hence, if  $\mathcal{P}^{(n)}$  in the previous section is assumed to be universal for all  $n$ , "sufficient" implies "summarizing" and "totally sufficient with  $\Sigma$ -structure" implies "summarizing with  $\Sigma$ -structure". Hence from the diagram it appears that "minimal totally sufficient" implies everything but minimal sufficient. Finally, if  $X_1, X_2, \dots$  are all assumed to be independent and at the same time  $\mathcal{P}^{(n)}$  assumed to be universal for all  $n$ , "minimal sufficient" implies everything else.



### References

- Bahadur, R.R. (1954): Sufficiency and statistical decision functions. Ann. Math. Statist., 25, 423-462.
- Barndorff-Nielsen, O. (1973): On M-ancillarity. Biometrika, 60, 447-455.
- Barndorff-Nielsen, O. and M. Skibinsky (1963): Adequate subfields and almost sufficiency. Appl. Math. Publ. No. 329, Brookhaven national laboratory, 39 pp.
- Fisher, R.A. (1920): A mathematical examination of the methods of determining the accuracy of an observation by the mean error, and by the mean square error. Monthly Notices R. Astronomical Soc., 80, 758-770. Reprinted (1950) in Fisher's Contributions to Mathematical Statistics. Wiley, New York.
- Freedman, D.A. (1962): Invariants under mixing which generalize de Finetti's theorem. Ann. Math. Statist., 33, 916-923.
- Lauritzen, S.L. (1972): Sufficiency and Time Series Analysis. Preprint No. 11. Inst. of Math. Statistics, University of Copenhagen.
- Lauritzen, S.L. (1974): Sufficiency, prediction and extreme models. Scand. J. Statist. 1. (to appear.)
- Martin-Löf, P. (1973): Repetitive structures. Inst. of Math. Statist. Univ. of Stockholm.
- Skibinsky, M. (1967): Adequate subfields and sufficiency. Ann. Math. Statist., 38, 155-161.

## TECHNICAL REPORTS

OFFICE OF NAVAL RESEARCH CONTRACT N00014-67-A-0112-0030 (NR-042-034)

1. "Confidence Limits for the Expected Value of an Arbitrary Bounded Random Variable with a Continuous Distribution Function," T. W. Anderson, October 1, 1969.
2. "Efficient Estimation of Regression Coefficients in Time Series," T. W. Anderson, October 1, 1970.
3. "Determining the Appropriate Sample Size for Confidence Limits for a Proportion," T. W. Anderson and H. Burstein, October 15, 1970.
4. "Some General Results on Time-Ordered Classification," D. V. Hinkley, July 30, 1971.
5. "Tests for Randomness of Directions against Equatorial and Bimodal Alternatives," T. W. Anderson and M. A. Stephens, August 30, 1971.
6. "Estimation of Covariance Matrices with Linear Structure and Moving Average Processes of Finite Order," T. W. Anderson, October 29, 1971.
7. "The Stationarity of an Estimated Autoregressive Process," T. W. Anderson, November 15, 1971.
8. "On the Inverse of Some Covariance Matrices of Toeplitz Type," Raul Pedro Mentz, July 12, 1972.
9. "An Asymptotic Expansion of the Distribution of "Studentized" Classification Statistics," T. W. Anderson, September 10, 1972.
10. "Asymptotic Evaluation of the Probabilities of Misclassification by Linear Discriminant Functions," T. W. Anderson, September 28, 1972.
11. "Population Mixing Models and Clustering Algorithms," Stanley L. Sclove, February 1, 1973.
12. "Asymptotic Properties and Computation of Maximum Likelihood Estimates in the Mixed Model of the Analysis of Variance," John James Miller, November 21, 1973.
13. "Maximum Likelihood Estimation in the Birth-and-Death Process," Niels Keiding, November 28, 1973.
14. "Random Orthogonal Set Functions and Stochastic Models for the Gravity Potential of the Earth," Steffen L. Lauritzen, December 27, 1973.
15. "Maximum Likelihood Estimation of Parameters of an Autoregressive Process with Moving Average Residuals and Other Covariance Matrices with Linear Structure," T. W. Anderson, December, 1973.
16. "Note on a Case-Study in Box-Jenkins Seasonal Forecasting of Time Series," Steffen L. Lauritzen, April, 1974.

TECHNICAL REPORTS

OFFICE OF NAVAL RESEARCH CONTRACT N00014-67-A-0112-0030 (NR-042-034)

17. "General Exponential Models for Discrete Observations," Steffen L. Lauritzen, May, 1974.
18. "On the Interrelationships Among Sufficiency, Total Sufficiency and Some Related Concepts," Steffen L. Lauritzen, June, 1974.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE  |                       | READ INSTRUCTIONS<br>BEFORE COMPLETING FORM                                 |
|--|-----------------------|---|
| 1. REPORT NUMBER<br>Technical Report No. 18  | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER   |
| 4. TITLE (and Subtitle)<br>On the Interrelationships Among Sufficiency, Total Sufficiency and Some Related Concepts.   |                       | 5. TYPE OF REPORT & PERIOD COVERED<br>Technical Report                      |
|  |                       | 6. PERFORMING ORG. REPORT NUMBER  |
| 7. AUTHOR(s)<br>Steffen L. Lauritzen   |                       | 8. CONTRACT OR GRANT NUMBER(s)<br>N00014-67-A-0112-0030                     |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS<br>Department of Statistics<br>Stanford University<br>Stanford, California 94305   |                       | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS<br>(NR-042-034) |
| 11. CONTROLLING OFFICE NAME AND ADDRESS<br>Office of Naval Research<br>Statistics and Probability Program Code 436<br>Arlington, Virginia 22217  |                       | 12. REPORT DATE   |
|  |                       | 13. NUMBER OF PAGES<br>15   |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)  |                       | 15. SECURITY CLASS. (of this report)<br>Unclassified                        |
|  |                       | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE                                  |
| 16. DISTRIBUTION STATEMENT (of this Report)<br><br>Reproduction in whole or in part is permitted for any purpose of the United States Government. Distribution is unlimited  |                       |   |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)   |                       |   |
| 18. SUPPLEMENTARY NOTES  |                       |   |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number)<br><br>adequacy, conditional independence, S-structure, statistical inference, stochastic processes, summarizing statistics, total sufficiency, transitivity, universality.   |                       |   |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)<br><br>As noted by various authors, the notion of sufficiency is too weak for problems in connection with statistical inference in stochastic processes. Various attempts have been made to impose extra conditions and in the present paper we shall discuss a few of these, with the purpose of discovering in which sense the concepts so defined differ and in which sense they are alike. |                       |   |